

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
**WO 02/087424 A2**

(51) International Patent Classification<sup>7</sup>: **A61B**

(21) International Application Number: PCT/US02/13890

(22) International Filing Date: 2 May 2002 (02.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/288,208 2 May 2001 (02.05.2001) US

(71) Applicants and

(72) Inventors: **LOW, Philip, Stewart** [US/US]; 5850 Farm Ridge Road, West Lafayette, IN 47906 (US). **TURK, Mary, Jo** [US/US]; 1233 York Avenue, Apt. 13E, New York, NY 10026 (US).

(74) Agent: **LAMMERT, Steven, R.**; Barnes & Thornburg, 11 South Meridian Street, Indianapolis, IN 46204 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TREATMENT AND DIAGNOSIS OF MACROPHAGE MEDIATED DISEASE

(57) Abstract: The invention relates to a method of treating or monitoring/diagnosing a disease state mediated by activated macrophages. The method comprises the step of administering to a patient suffering from a macrophage mediated disease state an effective amount of a composition comprising a conjugate or complex of the general formula:  $A_b-X$ , where the group  $A_b$  comprises a ligand capable of binding to activated macrophages, and when the conjugate is being used for treatment of the disease state, the group X comprises an immunogen, a cytotoxin, or a compound capable of altering macrophage function, and when the conjugate is being used for monitoring/diagnosing the disease state, X comprises an imaging agent. The method is useful for treating a patient suffering from a disease selected from the group consisting of rheumatoid arthritis, ulcerative colitis, Crohn's disease, inflammation, infections, osteomyelitis, atherosclerosis, organ transplant rejection, pulmonary fibrosis, sarcoidosis, and systemic sclerosis.

## TREATMENT AND DIAGNOSIS OF MACROPHAGE MEDIATED DISEASE

## FIELD OF THE INVENTION

This invention relates to methods for treating and monitoring disease states mediated by activated macrophages. More particularly, ligands that bind to activated macrophages are complexed with an imaging agent, or an immunogen, a cytotoxin or an agent for altering macrophage function for administration to a diseased host for diagnosis and/or treatment of macrophage mediated disease.

## 10 BACKGROUND AND SUMMARY OF THE INVENTION

The mammalian immune system provides a means for the recognition and elimination of foreign pathogens. While the immune system normally provides a line of defense against foreign pathogens, there are many instances where the immune response itself is involved in the progression of disease. Exemplary of diseases caused or worsened by the host's own immune response are autoimmune diseases such as multiple sclerosis, lupus erythematosus, psoriasis, pulmonary fibrosis, and rheumatoid arthritis and diseases in which the immune response contributes to pathogenesis such as atherosclerosis, inflammatory diseases, osteomyelitis, ulcerative colitis, Crohn's disease, and graft versus host disease often resulting in organ transplant rejection.

Macrophages are generally the first cells to encounter foreign pathogens, and accordingly, they play an important role in the immune response. However, activated macrophages can contribute to the pathophysiology of disease in some instances. Activated macrophages nonspecifically engulf and kill foreign pathogens within the macrophage by hydrolytic and oxidative attack resulting in degradation of the pathogen. Peptides from degraded proteins are displayed on the macrophage cell surface where they can be recognized by T cells, and they can directly interact with antibodies on the B cell surface, resulting in T and B cell activation and further stimulation of the immune response.

30 Rheumatoid arthritis (RA) is a systemic disease characterized by chronic inflammatory synovitis, usually involving peripheral joints. The synovial inflammation causes cartilage deterioration and bone erosion with consequent destruction of joint integrity. Rheumatoid factors, which are autoantibodies reactive

with the Fc region of IgG, are found in more than two-thirds of patients with RA indicating that RA has an autoimmune component.

RA is seen throughout the world in as much as 2% of the population, with 80% of RA patients developing the disease between the ages of 35 and 50. The clinical manifestations of RA include pain, swelling, and tenderness in the joints resulting in limitation of motion, weakness, fatigue, and weight loss. RA is a systemic disease and, consequently, has extra-articular manifestations, especially in patients with high titers of rheumatoid factors. These symptoms include rheumatoid nodules with an inner zone of necrotic material, a mid-zone of macrophages, and an outer zone of granulated tissue, muscle atrophy, osteoporosis, pulmonary fibrosis, and rheumatoid vasculitis which may result in cutaneous ulceration, digital gangrene, or neurovascular disease.

Rheumatoid synovitis, characteristic of RA, results in an increase in the number of synovial lining cells, hyperplasia and hypertrophy of the synovial lining cells, microvascular injury, edema, and infiltration of cells such as T cells, macrophages, and dendritic cells. The rheumatoid synovium is characterized by the presence of secreted products of immune cells such as factors secreted by T lymphocytes including IL-2, IFN- $\delta$ , IL-6, IL-10, GM-CSF and TGF $\alpha$  and  $\beta$  and factors secreted by activated macrophages including IL-1, IL-6, IL-8, IL-10, GM-CSF, macrophage CSF, and TGF $\beta$ . The production of these cytokines appears to account for much of the pathology of RA including inflammation of the synovium, synovial cell proliferation, cartilage and bone deterioration, and systemic symptoms of the disease.

RA may be treated using various therapies including physical therapy, rest, and splinting. Therapeutic agents are also used for the treatment of RA including aspirin and nonsteroidal anti-inflammatory drugs to control local inflammation. However, these agents have a minimal effect on the progression of the disease and are associated with toxic side effects. Disease-modifying anti-rheumatic drugs, such as  $\alpha$ -penicillamine and sulfasalazine, are also used to treat RA, but the benefit from these drugs is delayed for weeks or months and these drugs have toxic side effects. Immunosuppressive and cytotoxic drugs suppress symptoms of RA in some patients, but are associated with toxicity. Intra-articular glucocorticoids have also been used, but provide only transient relief. Accordingly, there is a need for the development of

new therapies with reduced toxicity that are efficacious for the treatment of RA and other diseases caused or worsened by activated macrophages.

The folate receptor (FR) is a 38 KDa GPI-anchored protein that binds the vitamin folic acid with high affinity ( $< 1$  nM). Following receptor binding, rapid  
5 endocytosis delivers the vitamin into the cell, where it is unloaded in an endosomal compartment at low pH. Importantly, covalent conjugation of small molecules, proteins, and even liposomes to folic acid does not alter the vitamin's ability to bind the folate receptor, and therefore, folate-drug conjugates can readily enter cells by receptor-mediated endocytosis.

10 Because most cells use an unrelated reduced folate carrier (RFC) to acquire the necessary folic acid, expression of the folate receptor is restricted to a few cell types. With the exception of kidney and placenta, normal tissues express low or nondetectable levels of FR. However, many malignant tissues, including ovarian, breast, bronchial, and brain cancers express significantly elevated levels of the  
15 receptor. In fact, it is estimated that 95% of all ovarian carcinomas overexpress the folate receptor. It has recently been reported that FR<sub>B</sub>, the nonepithelial isoform of the folate receptor, is expressed on activated (but not resting) synovial macrophages. Thus, Applicants have attempted to utilize folate-linked compounds potentially capable of altering the function of activated macrophages, to treat macrophage-  
20 mediated disease states. For example, Applicants have found that folate-linked immunogens can be used to redirect the host immune response in arthritic animals to activated macrophages at the site of inflammation to deplete macrophages and reduce arthritic inflammation.

Scintigraphic imaging agents are a million times more sensitive than  
25 magnetic resonance imaging (MRI) contrast agents, and their selectivity can be enhanced by their targeting to lesion-specific cell markers. Indeed, the radioisotope <sup>99m</sup>Tc has been delivered to arthritic tissues using nonspecific IgG, anti-CD4 antibodies, CD11b/CD14-glycolipopeptide ligands, and E-selectin binding peptides. Preclinical studies with such radioimaging agents have clearly emphasized the value  
30 of imaging arthritic tissues *in-vivo*, however, the selectivity of the current imaging agents is not yet optimal, and none of the present compounds is targeted exclusively to activated macrophages. In view of the emergence of folate receptor activity during



macrophage activation, Applicants have undertaken to determine whether a folate-targeted  $^{99m}\text{Tc}$  imaging agent might be used to image arthritic lesions *in vivo*.

To determine whether expression of this high affinity FR might be exploited to selectively target drugs to activated macrophages at sites of inflammation, folic acid has been conjugated to a  $^{99m}\text{Tc}$  chelator, and its distribution evaluated in both normal and diseased tissues of rats with adjuvant-induced arthritis. The folate-linked  $^{99m}\text{Tc}$  chelate complex, termed EC20, was indeed found to concentrate in the arthritic extremities of diseased rats, but not in the joints of healthy rats. The intensity of the gamma scintigraphic images of affected tissues was found to be greatly reduced in the presence of excess competing folic acid. Furthermore, liver and spleen of arthritic animals also showed enhanced uptake of EC20 and increased levels of FR, confirming that systemic activation of macrophages accompanies adjuvant-induced arthritis. Depletion of macrophages from arthritic animals reduced tissue FR content and concomitantly abolished uptake of EC20. Furthermore, Kupffer cells isolated from rats with adjuvant-induced arthritis exhibited a significantly higher binding capacity for folate conjugates than Kupffer cells from healthy rats. Thus, Applicants have found that EC20 is useful for assaying the participation of activated macrophages in inflammatory pathologies such as rheumatoid arthritis.

The present invention is directed to a method for treating and monitoring disease states mediated by activated macrophages. In accordance with one embodiment of the present invention, disease states mediated by activated macrophages are treated by redirecting host immune responses to activated macrophages or by altering the function of activated macrophages or by direct killing of activated macrophages. In one aspect of the invention, to promote killing of activated macrophages, ligands that bind specifically to activated macrophages are conjugated with an immunogen to redirect host immune responses to the activated macrophage population, or they are conjugated to a cytotoxin for direct killing of macrophages. Ligands that can be used in the conjugates of the present invention include those that bind to receptors expressed specifically on activated macrophages, such as the folate receptor, or ligands such as monoclonal antibodies directed to cell surface markers specifically expressed on activated macrophages. In another aspect of the invention ligands that bind specifically to activated macrophages are

conjugated with an imaging agent; the conjugate is administered to a patient for diagnosing and monitoring the progression of diseases mediated by activated macrophages.

5 In one embodiment, a method of treating or monitoring/diagnosing a disease state mediated by activated macrophages is provided. The method comprises the step of administering to a patient suffering from a macrophage mediated disease state an effective amount of a composition comprising a conjugate or complex of the general formula  $A_b-X$ , where the group  $A_b$  comprises a ligand capable of binding to activated macrophages, and when the conjugate is being used for treatment of the  
10 disease state, the group X comprises an immunogen, a cytotoxin, or a compound capable of altering macrophage function, and when the conjugate is being used for monitoring/diagnosing the disease state, X comprises an imaging agent.

#### DETAILED DESCRIPTION OF THE INVENTION

15 Methods are provided in accordance with the present invention for either treating or monitoring/diagnosing a disease state mediated by activated macrophages. Exemplary of diseases known to be mediated by activated macrophages include rheumatoid arthritis, ulcerative colitis, Crohn's disease, psoriasis, osteomyelitis, multiple sclerosis, atherosclerosis, pulmonary fibrosis, sarcoidosis, systemic sclerosis, organ transplant rejection (GVHD) and chronic  
20 inflammations. Such disease states can be monitored by first administering to a patient suffering from such disease state an effective amount of a composition comprising a conjugate of the general formula  $A_b-X$  wherein the group  $A_b$  comprises a ligand capable of binding to activated macrophages, and the group X comprises an  
25 imaging agent and thereafter scanning the patient with an imaging device capable of detecting the imaging agent. Macrophage mediated disease states can be treated in accordance with this invention by administering an effective amount of a composition of the above formula wherein  $A_b$  comprises a ligand capable of binding to an activated macrophage and wherein the group X comprises an immunogen, a  
30 cytotoxin, or a cytokine capable of altering macrophage function. Such macrophage targeting conjugates, when administered to a patient suffering from an activated macrophage mediated disease state, work to concentrate and associate the conjugated cytotoxin, immunogen, or cytokine with the population of activated macrophages to

kill the activated macrophages or alter macrophage function. Elimination or deactivation of the activated macrophage population works to stop or reduce the activated macrophage mediated pathogenesis characteristic of the disease state being treated. The conjugate is typically administered parenterally as a composition comprising the conjugate and a pharmaceutically acceptable carrier therefor. Conjugate administration is typically continued until symptoms of the disease state are reduced or eliminated.

In one embodiment of the invention activated macrophage mediated disease states are monitored or diagnosed in a patient by administering a conjugate  $A_b$ -X wherein  $A_b$  comprises a ligand capable of binding to activated macrophages and X comprises an imaging agent and thereafter scanning the patient with an imaging device capable of detecting localized concentration of the imaging agent. The imaging or diagnostic conjugates are, similar to those therapeutic conjugates outlined above, typically administered as a diagnostic composition comprising a conjugate and a pharmaceutically acceptable carrier. The composition is typically formulated for parenteral administration and is administered to the patient in an amount effective to enable imaging of the locale of activated macrophage populations. The nature of the imaging agent component of the conjugate is dictated by the imaging methodology. Thus, for example, the imaging agent can comprise a chelating moiety and a metal cation, for example, a radionuclide or a nuclear resonance imaging contrast agent, such as gadolinium. Typically the activated macrophage targeted imaging agent is administered to a patient, and following a period of time to allow delivery and concentration of the imaging agent in the activated macrophage cell populations, the patient is subjected to the imaging procedure and imaging is enabled by the targeted imaging agent.

The method of the present invention can be used for both human clinical medicine and veterinary applications. Thus, the host animals afflicted with the activated macrophage mediated disease state can be humans, or in the case of veterinary applications, they can be laboratory, agricultural, domestic or wild animals. The conjugates administered in accordance with the methods of this invention are preferably administered parenterally to the animal or patient suffering from the disease state, for example, intradermally, subcutaneously, intramuscularly, intraperitoneally, or intravenously. Alternatively, the conjugates can be administered

to the animal or patient by other medically useful procedures and effective doses can be administered in standard or prolonged release dosage forms, such as a slow pump. The therapeutic method of the present invention may be used alone or in combination with other therapeutic methods recognized for the treatment of macrophage mediated  
5 disease states.

In the ligand conjugates of the general formula  $A_b-X$  in accordance with the present invention, the group  $A_b$  is a ligand capable of binding to activated macrophages. Any of a wide number of macrophage binding moieties can be employed. Acceptable ligands include particularly folate receptor binding ligands and  
10 antibodies or antibody fragments capable of recognizing and specifically binding to surface moieties uniquely or preferentially expressed or presented in/on activated macrophages. In one embodiment the activated macrophage binding ligand is folic acid, a folic acid analog or other folate receptor binding molecules. Activated macrophages express a 38 LD GPI-anchored folate receptor that binds folate and  
15 folate-derivatized compounds with subnanomolar affinity (*i.e.*,  $< 1nM$ ). In another embodiment the activated macrophage binding ligand is a specific monoclonal or polyclonal antibody or Fab or scFv (*i.e.*, a single chain variable region) fragments of antibodies capable of specific binding to activated macrophages.

The activated macrophage targeted conjugates used for diagnosing and  
20 monitoring disease states mediated by activated macrophages in accordance with this invention are formed to target and, thus, to concentrate an imaging agent at the site of activated macrophage populations in the diseased patient. In such conjugates of the formula  $A_b-X$ ,  $A_b$  is a ligand capable of binding to activated macrophages and the group X comprises an imaging agent. In one embodiment the imaging agent  
25 comprises a chelating agent and a metal cation, typically either a radionuclide or a nuclear magnetic resonance imaging enhancer or contrast agent, such as gadolinium. Such conjugates wherein the group  $A_b$  is folic acid, a folic acid analog, or another folic acid receptor binding ligand are described in detail in U.S. Patent No. 5,688,488, the specification of which is incorporated herein by reference. That patent, as well as  
30 related U.S. Patents Nos. 5,416,016 and 5,108,921, each incorporated herein by reference, describe methods and examples for preparing chelate conjugates useful in accordance with the present invention. The present macrophage targeted imaging agents can be prepared and used following general protocols described in those earlier



patents. The present diagnostic method, however, is based in part on the discovery that folate targeted conjugates can be used to concentrate conjugated imaging entities in and at activated macrophage populations enabling monitoring and diagnosis of disease states characterized by concentration of activated macrophages at the site of disease.

In accordance with one embodiment of the present invention there is provided a method of treating disease states mediated by activated macrophages by administering to a patient suffering from such disease state an effective amount of a composition comprising a conjugate of the general formula  $A_b-X$  wherein  $A_b$  is as defined above and the group X comprises a cytotoxin, an immunogen, or a compound capable of altering macrophage function. Exemplary of cytotoxic moieties useful for forming conjugates for use in accordance with the present method include clodronate, anthrax, Pseudomonas exotoxin, typically modified so that these cytotoxic moieties do not bind to normal cells, and other toxins or cytotoxic agents including art-recognized chemotherapeutic agents such as adrenocorticoids, alkylating agents, antiandrogens, antiestrogens, androgens, estrogens, antimetabolites such as cytosine arabinoside, purine analogs, pyrimidine analogs, and methotrexate, busulfan, carboplatin, chlorambucil, cisplatin and other platinum compounds, tamoxiphen, taxol, cyclophosphamide, plant alkaloids, prednisone, hydroxyurea, teniposide, and bleomycin, nitrogen mustards, nitrosureas, vincristine, vinblastine, inflammatory and proinflammatory agents, and the like. Such toxins or cytotoxic components can be directly conjugated to the activated macrophage binding moiety, for example, folate or other folate receptor binding ligands, or they can be formulated in liposomes which themselves are targeted as conjugates of macrophage binding entities typically by covalent linkages to component phospholipids. Similarly, when the group X comprises a compound capable of altering a macrophage function, for example, a cytokine such as IL-10 or IL-11, the cytokine can be covalently linked to the targeting moiety  $A_b$ , for example, a folate receptor binding ligand or an antibody or antibody fragment directly, or the macrophage function altering cytokine can be encapsulated in a liposome which is itself targeted to activated macrophages by pendent macrophage targeting entities  $A_b$  covalently linked to one or more phospholipid liposome components.

In another embodiment the ligand-immunogen conjugates can be administered in combination with a cytotoxic compound. The compounds listed in the preceding paragraph are among the compounds suitable for this purpose.

In another method of treatment embodiment of the present invention  
5 the group X in the activated macrophage targeted conjugate  $A_b$ -X, comprises an immunogen, the ligand-immunogen conjugates being effective to "label" the population of activated macrophages responsible for disease pathogenesis in the patient suffering from the disease for specific elimination by an endogenous immune response or by co-administered antibodies. The use of ligand-immunogen conjugates  
10 in the method of treatment in accordance with this invention works to enhance an immune response-mediated elimination of the activated macrophage population. Such can be effected through an endogenous immune response or by a passive immune response effected by co-administered antibodies. The endogenous immune response may include a humoral response, a cell-mediated immune response, and any  
15 other immune response endogenous to the host animal, including complement-mediated cell lysis, antibody-dependent cell-mediated cytotoxicity (ADCC), antibody opsonization leading to phagocytosis, clustering of receptors upon antibody binding resulting in signaling of apoptosis, antiproliferation, or differentiation, and direct immune cell recognition of the delivered antigen/hapten. It is also contemplated that  
20 the endogenous immune response will employ the secretion of cytokines that regulate such processes as the multiplication and migration of immune cells. The endogenous immune response may include the participation of such immune cell types as B cells, T cells, including helper and cytotoxic T cells, macrophages, natural killer cells, neutrophils, LAK cells, and the like.

25 In another embodiment, the ligand-immunogen conjugate can be internalized and the immunogen can be degraded and presented on the macrophage cell surface for recognition by immune cells to elicit an immune response directed against macrophages presenting the degraded immunogen.

The humoral response may be a response induced by such processes as  
30 normally scheduled vaccination, or active immunization with a natural antigen or an unnatural antigen or hapten, *e.g.*, fluorescein isothiocyanate (FITC), with the unnatural antigen inducing a novel immunity. Active immunization involves multiple injections of the unnatural antigen or hapten scheduled outside of a normal

vaccination regimen to induce the novel immunity. The humoral response may also result from an innate immunity where the host animal has a natural preexisting immunity, such as an immunity to  $\alpha$ -galactosyl groups. Alternatively, a passive immunity may be established by administering antibodies to the host animal such as

5 natural antibodies collected from serum or monoclonal antibodies that may or may not be genetically engineered antibodies, including humanized antibodies. The utilization of a particular amount of an antibody reagent to develop a passive immunity, and the use of a ligand-immunogen conjugate wherein the passively administered antibodies are directed to the immunogen, would provide the advantage

10 of a standard set of reagents to be used in cases where a patient's preexisting antibody titer to other potential antigens is not therapeutically useful. The passively administered antibodies may be "co-administered" with the ligand-immunogen conjugate, and co-administration is defined as administration of antibodies at a time prior to, at the same time as, or at a time following administration of the ligand-

15 immunogen conjugate. It is contemplated that the preexisting antibodies, induced antibodies, or passively administered antibodies will be redirected to the activated macrophages by preferential binding of the ligand-immunogen conjugates to the activated macrophage cell populations, and such pathogenic cells are killed by complement-mediated lysis, ADCC, antibody-dependent phagocytosis, or antibody

20 clustering of receptors. The cytotoxic process may also involve other types of immune responses, such as cell-mediated immunity, as well as secondary responses that arise when the attracted antigen-presenting cells phagocytose the activated macrophages and present antigens of such cells to the immune system for elimination of other activated macrophages

25 presenting such antigens.

Acceptable immunogens for use in preparing the conjugates used in the method of treatment of the present invention are immunogens that are capable of eliciting antibody production in a host animal or that have previously elicited antibody production in a host animal, resulting in a preexisting immunity, or that

30 constitute part of the innate immune system. Alternatively, antibodies directed against the immunogen may be administered to the host animal to establish a passive immunity. Suitable immunogens for use in the invention include antigens or antigenic peptides against which a preexisting immunity has developed via normally

scheduled vaccinations or prior natural exposure to such agents such as polio virus, tetanus, typhus, rubella, measles, mumps, pertussis, tuberculosis and influenza antigens and  $\alpha$ -galactosyl groups. In such cases, the ligand-immunogen conjugates will be used to redirect a previously acquired humoral or cellular immunity to a  
5 population of activated macrophages in the host animal for elimination of such cells. Other suitable immunogens include antigens or antigenic peptides to which the host animal has developed a novel immunity through immunization against an unnatural antigen or hapten, for example, fluorescein isothiocyanate (FITC) or dinitrophenyl and antigens against which an innate immunity exists, for example, super antigens  
10 and muramyl dipeptide. It is also contemplated that MHC I restricted peptides could be linked to the ligand for use in redirecting cellular immunity to macrophages and eliciting T cell killing of macrophages.

The macrophage binding ligands and immunogens, cytotoxic agents, cytokines or imaging agents, as the case may be in forming conjugates for use in  
15 accordance with the present invention, may be conjugated by using any art-recognized method for forming a complex. This can include covalent, ionic, or hydrogen bonding of the ligand to the immunogen, either directly or indirectly via a linking group such as a divalent linker. The conjugate is typically formed by covalent bonding of the ligand to the targeted entity through the formation of amide, ester or  
20 imino bonds between acid, aldehyde, hydroxy, amino, or hydrazo groups on the respective components of the complex. Alternatively, as mentioned above, the ligand complex can be one comprising a liposome wherein the targeted entity (that is, the imaging agent, or the immunogen, cytotoxic agent or macrophage function altering agent) is contained within a liposome which is itself covalently linked to the activated  
25 macrophage binding ligand.

In one embodiment of the invention the ligand is folic acid, an analog of folic acid, or any other folate receptor binding molecule, and the folate ligand is conjugated to the targeted entity by a procedure that utilizes trifluoroacetic anhydride to prepare  $\gamma$ -esters of folic acid via a pteroyl azide intermediate. This procedure  
30 results in the synthesis of a folate ligand, conjugated to the targeted entity only through the  $\gamma$ -carboxy group of the glutamic acid groups of folate. Alternatively, folic acid analogs can be coupled through the  $\alpha$ -carboxy moiety of the glutamic acid group or both the  $\alpha$  and  $\gamma$  carboxylic acid entities.



The conjugates used in accordance with this invention of the formula  $A_b-X$  are used in one aspect of this invention to formulate therapeutic or diagnostic compositions comprising effective amounts of the conjugate and an acceptable carrier therefor. Typically such compositions are formulated for parenteral use. The amount  
5 of the conjugate effective for use in accordance with the invention depends on many parameters, including the nature of the disease being treated or diagnosed, the molecular weight of the conjugate, its route of administration and its tissue distribution, and the possibility of co-usage of other therapeutic or diagnostic agents. The effective amount to be administered to a patient is typically based on body  
10 surface area, patient weight and physician assessment of patient condition. An effective amount can range from about 1 ng/kg to about 1 mg/kg, more typically from about 1  $\mu$ g/kg to about 500  $\mu$ g/kg, and most typically from about 1  $\mu$ g/kg to about 100  $\mu$ g/kg.

When used for monitoring or diagnosis, imaging procedures are  
15 typically carried out about 1 to about 6 hours post administration of the activated macrophage targeted imaging agent.

Any effective regimen for administering the ligand conjugates can be used. For example, the ligand conjugates can be administered as single doses, or they can be divided and administered as a multiple-dose daily regimen. Further, a  
20 staggered regimen, for example, one to three days per week can be used as an alternative to daily treatment, and for the purpose of defining this invention such an intermittent or staggered daily regimen is considered to be equivalent to every day treatment and within the scope of this invention. In one embodiment of the invention the patient is treated with multiple injections of the ligand conjugate wherein the  
25 targeted entity is an immunogen or a cytotoxic agent to eliminate the population of pathogenic activated macrophages. In one embodiment, the patient is treated, for example, injected multiple times with the ligand conjugate at, for example, at 12-72 hour intervals or at 48-72 hour intervals. Additional injections of the ligand conjugate can be administered to the patient at intervals of days or months after the initial  
30 injections, and the additional injections prevent recurrence of disease. Alternatively, the ligand conjugates may be administered prophylactically to prevent the occurrence of disease in patients known to be disposed to development of activated macrophage mediated disease states. In one embodiment of the invention more than one type of

ligand conjugate can be used, for example, the host animal may be pre-immunized with fluorescein isothiocyanate and dinitrophenyl and subsequently treated with fluorescein isothiocyanate and dinitrophenyl linked to the same or different activated macrophage targeting ligands in a co-dosing protocol.

5                   The ligand conjugates are administered in accordance with this invention parenterally and most typically by intraperitoneal injections, subcutaneous injections, intramuscular injections, intravenous injections or intrathecal injections. The ligand conjugates can also be delivered to a patient using an osmotic pump. Examples of parenteral dosage forms include aqueous solutions of the conjugate, for  
10                   example, solution in isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carriers such as alcohols, glycols, esters and amides. The parenteral compositions for use in accordance with this invention can be in the form of a reconstitutable lyophilizate comprising the one or more doses of the ligand conjugate. In another aspect of the invention, the ligand conjugates can be  
15                   formulated as one of any of a number of prolonged release dosage forms known in the art such as, for example, the biodegradable carbohydrate matrices described in U.S. Patent Nos. 4,713,249; 5,266,333; and 5,417,982, the disclosures of which are incorporated herein by reference.

#### EXAMPLE 1

#### 20                   MATERIALS

                  EC20 (a folate-linked chelator <sup>99m</sup>Tc), EC28 (the same <sup>99m</sup>Tc chelate complex without folate), and folate-fluorescein isothiocyanate (folate-FITC) were gifts from Endocyte, Inc. (West Lafayette, IN). Heat-killed *Mycoplasma butericum* was purchased from BD Biosciences (Sparks, MD). Folic acid, light mineral oil,  
25                   clodronate, collagenase-A, and streptavidin-R-phycoerythrin were obtained from Sigma Chemical Co. (St. Louis, MO), and Dubelco's Modified Eagle Medium (DMEM) was from Gibco-BRL (Gathersberg, MD). <sup>3</sup>H-folic acid was obtained from American Radiolabeled Chemicals, Inc. (St. Louis, MO) and Microcon®-30  
30                   membranes were purchased from Millipore Corp. (Bedford, MA). RK-4-biotin and ED2-R-phycoerythrin antibodies were acquired from Bachem Biosciences, Inc. (Philadelphia, PA) and Accurate Chemical and Scientific Corp. (Westbury, NY), respectively.

-14-

## EXAMPLE 2

## ANIMAL MODEL OF ARTHRITIS

Arthritis was induced in 150-200g female Lewis rats (Charles River Laboratories, Inc., Wilmington, MA), n=4/dose group. Briefly, 0.5 mg of heat-killed  
5 *Mycoplasma butericum*, suspended in mineral oil (5 mg/ml), was injected on day 0 into the left hind foot of rats following anesthesia with ketamine and xylazine. Disease was allowed to progress for 21 days, and animals were weighed on a daily basis to ensure the status of their health. All treated animals developed arthritis, as evidenced by dramatic swelling in the injected paw, progressive swelling in all  
10 noninjected limbs due to the systemic progression of arthritis, and radiographic analysis of affected limbs. All rats were maintained on a folate-deficient diet (DYETS, Inc., Bethlehem, PA) for 3 weeks prior to administration of folate-FITC in order to lower serum folate levels to physiologically relevant concentrations. Control rats were also maintained on a folate-deficient diet but not induced to develop  
15 arthritis.

## EXAMPLE 3

## ELIMINATION OF ENDOGENOUS MACROPHAGES

Evaluation of macrophage independent uptake of the folate-linked imaging agent was accomplished by killing endogenous macrophages with liposomal  
20 clodronate. Liposomes were formed by rehydrating a thin film of egg phosphatidylcholine (60 mole %) and cholesterol (40 mole %) in an isotonic clodronate solution (250 mg/ml). Small unilamellar vesicles were then generated by extrusion of the liposomes ten times through a 100 nm polycarbonate membrane using a 10 ml thermobarrel extruder (Lipex Biomembranes, Vancouver, Canada).  
25 Unencapsulated clodronate was removed by dialysis through a Spectrapor 300,000 M<sub>r</sub>-cutoff cellulose acetate membrane (Spectrum Laboratories, Rancho Domingues, CA), and the clodronate concentration in the retained liposomes was determined as described in *J. Microencapsul.* 3(2) 109-14 (1986). Seventeen days following induction of the arthritis and three days prior to administration of the imaging agent  
30 (EC20), rats destined for macrophage depletion received a single intraperitoneal injection of clodronate liposomes containing 20 mg clodronate.

## EXAMPLE 4

## SCINTIGRAPHY AND BIODISTRIBUTION ANALYSIS

Twelve hours prior to administration of imaging agent, all animals received 5 ml of normal saline subcutaneously to ensure proper excretion of unbound  
5 imaging agent. Twenty-one days following induction of arthritis, rats (n = 3 per group) were injected intraperitoneally with 500  $\mu$ Ci (2.3 nmoles/rat) of either EC20 (folate + chelator), EC20 + 500-fold molar excess folic acid, or EC28 (no folate moiety). Four hours later, rats underwent either nuclear scintigraphic imaging or biodistribution analysis.

10 For scintigraphy, rats were anesthetized with ketamine and xylazine, and positioned in ventral recumbency on the image acquisition surface. Image acquisition was performed for one minute at a count rate of 50-75,000 counts per minute using a Technicare Omega 500 Sigma 410 Radioisotope Gamma Camera. Following acquisition of whole body images, radiation of the upper body (above the  
15 stifles) was blocked using 1/8" lead plates, and images of the posterior limbs were obtained. All data were analyzed using a Medasys™ MS-DOS-based computer equipped with Medasys™ Pinnacle software.

For biodistribution analysis, rats were euthanized by intraperitoneal injection of nebutal or pentobarbitol sodium. Liver, spleen, heart, lungs, intestine,  
20 and kidneys were then harvested and radiation in each tissue was determined by counting in a gamma counter (Packard BioScience Co., Meridian, CT).

## EXAMPLE 5

## ASSAY OF TISSUE FOLATE RECEPTOR LEVELS

Folate receptor levels in each tissue were determined follows. Briefly,  
25 tissues were homogenized and cell membranes were isolated by centrifugation. Membrane proteins were solubilized overnight, transferred into a Microcon®-30 filtration device, and incubated with 50 nM  $^3$ H-folic acid. A duplicate of each sample, used to determine non-specific binding, was also exposed to 50 nM  $^3$ H-folic acid, but in the presence of 1000-fold excess unlabeled folic acid. After unbound  $^3$ H-  
30 folic acid was washed through the membrane, membrane protein with bound  $^3$ H-folic acid was recovered and counted in a scintillation counter (Packard BioScience Co.) to determine the number of active folate receptors per gram of tissue.



-16-

## EXAMPLE 6

## IDENTIFICATION OF THE FOLATE RECEPTOR EXPRESSING CELL TYPE IN LIVER

Arthritic and healthy rats were first anesthetized with ketamine and xylazine, and then a midline incision was made, starting in the lower abdomen and extending through the thoracic cavity. A 24-gauge catheter was inserted into the hepatic vein, and a 24-gauge needle was inserted in the cardiac left ventricle to serve as an outlet for the perfusion fluid. Rats were then perfused by delivery of normal saline, followed by collagenase A solution (0.05% in Gey's balanced salt solution) through the catheter. Each solution was perfused for two minutes at a rate of 20 ml/minute. Immediately after perfusion, livers were removed and the membranous outer tissue was dissected away. The remaining gelatinous tissue was suspended in collagenase-A solution (0.025% in DMEM) and incubated at 37°C for two hours in the presence of 1  $\mu$ M folate-FITC or 1  $\mu$ M folate-FITC + 1 mM folic acid. Cells were then washed three times to removed unbound folate-FITC and immediately prepared for flow cytometry.

## EXAMPLE 7

## FLOW CYTOMETRY SAMPLE PREPARATION AND ANALYSIS

Liver cell preparations, which had been exposed to folate-FITC, were treated for 10 min at 4°C with ammonium chloride lysis buffer (150 mM  $\text{NH}_4\text{Cl}$ , 10 mM  $\text{KHCO}_3$ , 1 mM EDTA, pH 7.4) to lyse red blood cells. Following three washes with phosphate buffered saline, the remaining cells were incubated for 1 h at 4°C with either ED2 R-Phycoerythrin-labeled mouse anti-rat macrophage antibody, or RK-4 biotin-labeled mouse-anti rat granulocyte antibody. Cells were again washed two times, and those that had received the biotinylated primary antibody were further incubated with streptavidin-R-Phycoerythrin for 30 minutes. Following two final washes, cells were examined for FITC and phycoerythrin dual color staining on a FACScan Coulter XL flow cytometer.

## EXAMPLE 8

## IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-INDUCED ARTHRITIS

The protocol described in Example 2 for arthritis induction was followed. The efficacy of a folate-FITC conjugate (folate-fluorescein isothiocyanate

-17-

conjugate) against adjuvant-induced arthritis in rats was investigated. Each rat used in the experiment was immunized at the base of the tail with FITC-KLH (150 µg) to induce antibodies against FITC on days -38 and -10 before administration of *Mycoplasma butericum* (adjuvant) to induce arthritis. The immunization of FITC-KLH was done in combination with an adjuvant (*i.e.*, such as TiterMax Gold (150 µg), Alum (150 µg), or GPI-100 (150 µg) which are all adjuvants to induce antibodies against FITC as opposed to the adjuvant used to induce arthritis). The immunized animals were then injected on day 0 in the left foot pad with 0.5 mg of heat-killed *Mycoplasma butyricum* (adjuvant) to initiate development of arthritis. Then on days 1, 2, 3, 9, 11, and 14, post-adjuvant (*Mycoplasma butyricum*) injection, the rats were injected intraperitoneally with either saline (control rats) or 2000 nmoles/kg of folate-FITC (FF). Calipers were used to measure left and right foot dimensions daily. With reference to Fig. 9, those measurements are plotted for both the adjuvant-injected feet (top two curves) and the non-treated feet (bottom two curves). A sudden increase in swelling of the adjuvant-injected feet is due to influx of neutrophils which have no folate receptors. Consequently, the immunotherapy has no impact on this phase of paw swelling. However, after about 10 days, activated macrophages invade both injected feet and uninjected feet, causing bone degradation and further inflammation. These activated macrophages have functional folate receptors, and, as shown, they are eliminated or reduced by binding folate-hapten conjugates such as folate-FITC.

#### EXAMPLE 9

##### FOLATE-TARGETED IMAGING OF ARTHRITIC RATS

The protocols described in Examples 2 and 4 were followed. As noted above, activated but not resting macrophages express a receptor for the vitamin folic acid. To determine whether folate might be exploited to target <sup>99m</sup>Tc to sites of arthritic inflammation, EC20, a folate-linked chelator of <sup>99m</sup>Tc (see Fig. 8) was administered intraperitoneally to rats (n=5/group) and scintigraphic images were acquired with a gamma camera. Due to the rapid clearance of EC20, excellent contrast was obtained by at least four hours post-administration (Fig. 1). Importantly, whole body uptake was significantly more intense in arthritic rats compared to healthy rats, and this uptake was greatly reduced when EC20 was administered together with a saturating dose of free folic acid. This suggests that uptake by all tissues is primarily determined by a folic-specific receptor.

Intense organ uptake of EC20 prevented visualization of limbs in whole body images of the arthritic rats. However, images of posterior limbs could be easily acquired when mid and upper body radiation was shielded. With such shielding, arthritic limbs displayed much greater EC20 uptake than healthy  
5 extremities, and this uptake was completely eliminated in the presence of excess free folic acid (Fig. 1). Furthermore, the left rear foot of the arthritic animals, where inflammation was most severe, displayed greater uptake than the less severely affected right rear foot (Fig. 1).

From the whole body images, it could be concluded that abdominal  
10 organs were responsible for a majority of EC20 uptake in the arthritic animals. To confirm this assessment, liver, spleen and kidney were removed and imaged separately (Fig. 2). Livers of arthritic rats demonstrated the highest uptake of EC20, while livers of healthy rats displayed minimal uptake. Only those spleens taken from arthritic rats could be visualized. Free folic acid completely blocked EC20 uptake in  
15 liver and spleen, however, the free vitamin only partially decreased uptake by the kidney.

#### EXAMPLE 10

##### EFFECTS OF MACROPHAGE DEPLETION

The protocols described in Examples 2, 3, and 4 were followed, except  
20 that 0.25 mCi of EC20 was administered. In order to determine whether macrophages might be responsible for the uptake of EC20, resident macrophages were systemically eliminated from arthritic rats using a liposomal clodronate preparation (n=3 rats/group). By four days after clodronate treatment, evaluation of paw size revealed that clodronate-treated rats were significantly less inflamed than untreated rats (data  
25 not shown). To determine whether macrophage elimination would influence uptake of the folate-linked imaging agent, EC20 biodistribution analysis was then performed on the clodronate-treated rats and compared to the same analysis of both healthy rats and arthritic rats not treated with clodronate. As shown in Fig. 3 (black bars represent arthritic rats, light gray bars represent healthy rats, and dark gray bars represent  
30 arthritic rats depleted of macrophages by clodronate treatment), depletion of macrophages decreased liver uptake of EC20 ~20-fold in arthritic rats, while retention in the spleen and intestine was reduced by a factor of three. In most tissues, clodronate treatment depressed EC20 uptake even below those levels observed in

healthy rats, confirming the hypothesis that activated macrophages account for most of EC20 retention in normal tissues. In contrast, kidney uptake of EC20 was elevated in rats depleted of macrophages, most likely because the decreased internalization of EC20 by activated macrophages rendered more EC20 available for binding to kidney  
5 folate receptors.

#### EXAMPLE 11

##### FOLATE RECEPTOR-MEDIATED UPTAKE OF EC20 IN ARTHRITIC TISSUES

The protocols described in Examples 2 and 4 were followed. Two additional biodistribution studies were conducted to confirm that EC20 uptake by  
10 tissues of arthritic rats is mediated by the folate receptor (n=3 rats/group). First, the biodistribution of EC20 was examined in the presence (light gray bars) and absence (black bars) of a 500-fold excess of free folic acid. As seen in Fig. 4, almost complete elimination of EC20 uptake was observed in all tissues except kidney, indicating that binding was indeed mediated by a folate receptor. In fact, excess folic acid  
15 competitively reduced EC20 retention in liver, spleen, heart, lung, intestine and blood to near background levels (Fig. 4). Second, to confirm the role of folate in EC20-mediated targeting of the chelated  $^{99m}\text{Tc}$ , the biodistribution of the same complex lacking a folate moiety (EC28) was also examined (dark gray bars). As also  
20 displayed in Fig. 4, uptake of EC28 was negligible in all tissues except kidney, where retention of the non-targeted complex was similar to that of EC20 in the presence of competing folic acid.

#### EXAMPLE 12

##### FOLATE RECEPTOR EXPRESSION IN VARIOUS TISSUES OF ARTHRITIC RATS

25 The protocols described in Examples 2, 4, and 5 were used. The above results suggest that the folate receptor is responsible for tissue uptake of EC20. In order to confirm this, Applicants attempted to directly quantitate the folate binding protein in various rat tissues. Active folate receptor could be detected in each of the major organs examined, and FR levels were significantly increased in arthritic rats  
30 (Fig. 5; black bars = arthritic rats; light gray bars = healthy rats). Further, FR content correlated well with uptake of EC20 seen in the biodistribution studies. In fact, the FR assay revealed roughly equivalent levels of receptor in arthritic liver and spleen, in accordance with the similar uptake of EC20 by the same organs (Fig. 4).



-20-

Significantly, systemic elimination of macrophages by clodronate treatment (dark gray bars) lowered folate receptor levels in all arthritic tissues (Fig. 5), also in good agreement with the EC20 biodistribution analysis. Finally, the FR assay confirmed that neither induction of arthritis nor clodronate treatment alters the levels of FR in  
5 kidney or heart, where FR is not thought to be associated with activated macrophages.

## EXAMPLE 13

EXPRESSION OF A FUNCTIONAL FOLATE RECEPTOR ON LIVER  
MACROPHAGES OF ARTHRITIC RATS

The protocols described in Examples 2, 4, 6 and 7 were used. To  
10 further confirm that the elevated uptake of EC20 in livers of arthritic rats is due to a macrophage population, livers were resected from collagenase-perfused rats and their disaggregated cells examined for folate conjugate uptake, using folate-FITC as a fluorescent marker for FR expression. By also labeling the same liver cell suspension with an antibody specific for rat liver macrophages, it was possible to demonstrate  
15 that macrophages are indeed the cell type that expresses elevated levels of folate receptor in arthritic animals (Fig 6). Thus, flow cytometric analysis revealed that 70% of the liver macrophages of arthritic rats bound folate-FITC compared to only 30% of the liver macrophages of healthy rats (Fig. 6). Further, the FITC intensity of the arthritic macrophages was higher than that of macrophages from healthy livers.  
20 Since binding of folate-FITC was suppressed in the presence of an excess of free folic acid (1 mM), we concluded that uptake of the folate conjugate by liver macrophages was mediated by the folate receptor.

Using an antibody specific for granulocytes, we also examined whether tissue infiltrating neutrophils might take up folate conjugates. Although very  
25 few neutrophils were found in the liver, those that were detected exhibited no binding capacity for folate-FITC (data not shown). Mac-1+ peripheral blood cells were also tested and similarly found to have no binding affinity for the folate-conjugate (data not shown). In fact, no peripheral blood cells sorted positive for FITC fluorescence, suggesting that only resident tissue macrophages (and clearly only a subpopulation of  
30 those) express FR in the liver.

Finally, to begin to explore whether activated macrophages might be targeted with folate-linked drugs in human patients, we obtained permission to examine the whole body images of the 28 suspected ovarian cancer patients enrolled

in a recently completed clinical trial of the gamma imaging agent,  $^{111}\text{In}$ -DTPA-Folate. As shown in Fig. 7, one patient displayed significant imaging agent uptake in the right knee, but not the left knee. Importantly, no other patients demonstrated any measurable joint uptake. Upon request, the attending physician contacted the  
5 anonymous patient and inquired whether she had been experiencing any chronic joint discomfort. The physician responded that the patient reported arthritis in the right knee.

## DISCUSSION

Activated macrophages are thought to be intimately involved in the  
10 pathogenesis of rheumatoid arthritis. Activated macrophages directly destroy joint tissue by secreting metalloproteinases and attracting/activating other immune cells by releasing cytokines. The quantitation of activated macrophages in joint tissues might be of diagnostic value, since activated macrophage content correlates well with articular destruction and poor disease prognosis in humans.

15 Gamma camera scintigraphy of rats receiving EC20 demonstrated that arthritic appendages are indeed illuminated by folate-targeted  $^{99\text{m}}\text{Tc}$ . In contrast, the legs and feet of healthy rats could not be visualized, demonstrating the selectivity of the imaging agent for arthritis applications. Although the intensities of internal organs also increased in adjuvant-induced arthritis, interference from such tissues did  
20 not appear to compromise the methodology, since gamma radiation from internal organs could be easily screened. The fact that excellent contrast can be obtained within one to two hours of EC20 injection further shows that imaging agent administration, gamma camera scintigraphy, and image analysis can be completed during the same examination.

25 Systemic activation of macrophages has been documented in rats with adjuvant-induced arthritis. Thus, it was important to establish the specific participation of macrophages in the elevated uptake of EC20, since a folate-targeted imaging agent had never previously been examined in arthritic animals. Three experiments were conducted for this purpose. First, clodronate-loaded liposomes  
30 were employed to systemically deplete macrophages from the treated rats. Not only were the resulting tissue FR levels greatly reduced, but uptake of EC20 in the macrophage-rich organs was also nearly eliminated, suggesting that resident macrophages can indeed account for both FR expression and EC20 retention in the

RES organs. Second, liver cells were disaggregated by collagenase treatment and individual cells were evaluated for folate conjugate uptake. As noted in Fig. 6, the vast majority of cells testing positive for folate conjugate uptake also sorted positive for the macrophage marker, ED2, confirming that FR is indeed present on the  
5 macrophages. Finally, because other immune and myelocytic cells are known to be elevated in tissues of rats with adjuvant-induced arthritis, it was conceivable that still another extravasating blood cell type might be involved in the uptake of EC20. However, neither liver-infiltrating granulocytes nor any blood cell in circulation displayed any capacity to bind folate-FITC. Thus, activated macrophages would  
10 seem to be the predominant cell type internalizing folate conjugates in the organs of arthritic rats.

It was surprising to find that up to 30% of the liver macrophages in healthy rats also expressed the folate receptor (Fig. 6). Since a functional folate receptor is not found on resting synovial macrophages, it is tempting to speculate that  
15 the folate-FITC binding fraction in the healthy rats might also constitute an activated population. Two observations may support this conjecture. First, activated macrophages are also found in healthy tissues following exposure to immune stimulants such as foreign antigens. Given the role of the liver in clearing foreign substances from the body, a low level of resident macrophage activation does not  
20 seem unreasonable. Second, the folate-FITC (and EC20) binding population of liver cells increased significantly upon induction of localized inflammation and systemic macrophage activation.

With the ability to exploit folate to deliver attached molecules to activated macrophages now established, folate-linked imaging agents will allow the  
25 early development or continued progression of rheumatoid arthritis to be assessed. Since graft versus host disease, multiple sclerosis, Crohn's disease, ulcerative colitis, psoriasis, osteomyelitis, and even atherosclerosis may also be caused/aggravated by activated macrophages, it is possible that the diagnosis/evaluation of these diseases could be aided by a folate-linked imaging/contrast agent. The avid folate conjugate  
30 uptake by activated macrophages in both arthritic joints and liver indicates that macrophages can be effectively targeted regardless of their anatomical location.

-23-

## EXAMPLE 14

IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-  
INDUCED ARTHRITIS

For the assay shown in Fig. 10, the protocol described in Example 8  
5 was followed except that 3000 nmoles/kg of folate-FITC was administered per day (3  
doses on days 1, 2, and 3) and folate-FITC was delivered using an osmotic pump  
implanted into the peritoneal cavity of the rat. Methotrexate (MTX) was administered  
at a dose of 0.15 mg by intraperitoneal injection one time per day on days 1, 8, and 15  
after adjuvant administration. MTX was used in place of folate-FITC for animals  
10 treated with MTX. The results for both the left (injected) and right (uninjected) paw  
are shown. The results show that folate-FITC (FF) inhibits adjuvant-induced arthritis  
as well as MTX.

## EXAMPLE 15

IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-  
INDUCED ARTHRITIS

15

The protocol described in Example 14 was followed except that only  
the right paw volume was measured and folate-FITC (FF) was administered at doses  
of 3000, 600, and 120 nmoles/kg (Fig. 11). Also, FF was administered at 3000  
nmoles/kg in either three doses as in Example 14 (indicated as "3 doses" in Fig. 11) or  
20 on days 1, 2, 3, 9, 11, and 14 as described in Example 8 (indicated as "FF3000" in  
Fig. 11). The results show that FF inhibits adjuvant-induced arthritis in the right paw  
of the arthritic rats (inflammation presumably appears in the uninjected right paw due  
to the systemic progression of arthritis), and that prolonged treatment with FF is more  
effective than 3 initial doses for treatment of adjuvant-induced arthritis.

25

## EXAMPLE 16

IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-  
INDUCED ARTHRITIS

The protocol described in Example 8 was followed except MTX was  
used at a dose of 0.15 mg (FF + low, early MTX, late MTX and lateFFMTX) to treat  
30 some animals in place of FF (see Fig. 12). Other animals were treated with 0.75 mg  
of MTX (FF + mid) or 1.5 mg of MTX (FF + high). For "early MTX" treatments, the  
rats were injected with MTX on days 1, 8, and 15 after arthritis induction. For "late  
MTX" treatments, the rats were injected with MTX on days 8 and 15 after adjuvant



administration. All measurements were of the uninjected right paw. The results show that folate-FITC (FF) in combination with MTX (early or late treatments and a low, high, or middle dose of MTX) inhibits adjuvant-induced arthritis better than FF alone.

#### EXAMPLE 17

#### 5 IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-INDUCED ARTHRITIS

For the results shown in Fig. 13, the protocol described in Example 8 was followed except that some animals were treated with MTX alone (0.15 mg) on days 1, 8, and 15 after arthritis induction or were treated with MTX (0.15 mg; days 1,  
10 8, and 15) in combination with FF as described in Example 14. The results show that the combination of FF and MTX inhibits adjuvant-induced arthritis to a greater extent than MTX or FF alone.

#### EXAMPLE 18

#### 15 IMMUNOTHERAPY MEDIATED PROTECTION AGAINST ADJUVANT-INDUCED ARTHRITIS

The protocol described in Example 8 was followed except that IL-10 (10,000 U; FF10) or IL-2 (3 µg/kg; FF2) was administered along with the treatments with FF (*i.e.*, the cytokines were administered by intraperitoneal injections on days 1, 2, 3, 9, 11, and 14 after adjuvant administration; see Fig. 14). The measurements  
20 made were measurements of the right, noninjected paw. The results show that either IL-10 or IL-2 prevent the inhibition of adjuvant-induced arthritis resulting from treatment with FF. All of the above immunotherapy results taken together indicate that folate-linked agents which are cytotoxic for macrophages can be used to treat macrophage-mediated disease states.

#### 25 EXAMPLE 19

#### FOLATE-TARGETED IMAGING OF ARTHRITIC RATS

For the assay shown in Fig. 15, the protocols were as described in Examples 2 and 4 except that some animals were treated with FF (2000 nmoles/kg; days 1, 2, 3, 9, 11, and 14) or MTX (0.15 mg; days 1, 8, and 15) as described in  
30 Examples 8 and 14, respectively. The results show that FF or MTX prevent EC20 uptake in all organs examined except the kidney. It is likely that EC20 uptake is reduced in most organs making more EC20 available for excretion through the kidney accounting for the increase in EC20 detected in kidney tissues.

## CLAIMS

1. A method of treating or monitoring/diagnosing a disease state mediated by activated macrophages, said method comprising the step of administering  
5 to a patient suffering from a macrophage mediated disease state an effective amount of a composition comprising a conjugate or complex of the general formula  
$$A_b-X$$
where the group  $A_b$  comprises a ligand capable of binding to activated macrophages and when the conjugate is being used for treatment of the disease state,  
10 the group X comprises an immunogen, a cytotoxin, or a compound capable of altering macrophage function and when the conjugate is being used for monitoring/diagnosing the disease state, X comprises an imaging agent.
2. The method of claim 1 wherein  $A_b$  comprises a folate receptor binding ligand.
- 15 3. The method of claim 1 wherein  $A_b$  comprises an antibody or an activated macrophage binding antibody fragment.
4. The method of claim 1 wherein the group X comprises an immunogen.
5. The method of claim 2 wherein the group X comprises an  
20 immunogen.
6. The method of claim 1 wherein the group X comprises a cytotoxin.
7. The method of claim 6 wherein the group further comprises a liposome.
- 25 8. The method of claim 2 wherein the group X comprises a cytotoxin.
9. The method of claim 8 wherein the group X further comprises a liposome.
10. The method of claim 1 wherein the group X comprises an  
30 imaging agent.
11. The method of claim 10 wherein the group X further comprises a liposome.

12. The method of claim 2 wherein the group X comprises an imaging agent.
13. The method of claim 12 wherein the group X further comprises a liposome.
- 5 14. The method of claim 10 wherein the imaging agent comprises a metal chelating moiety.
15. The method of claim 14 wherein the imaging agent further comprises a metal cation.
- 10 16. The method of claim 15 wherein the metal cation is a radionuclide.
17. The method of claim 15 wherein the metal cation is a nuclear magnetic resonance imaging enhancing agent.
18. The method of claim 1 wherein X comprises a cytokine capable of altering activated macrophage function.
- 15 19. The method of claim 1 wherein the patient is suffering from a disease state selected from the group consisting of rheumatoid arthritis, ulcerative colitis, Crohn's disease, inflammatory lesions, infections of the skin, osteomyelitis, atherosclerosis, organ transplant rejection, pulmonary fibrosis, sarcoidosis, systemic sclerosis, and any chronic inflammation.




Fig. 1






Fig. 2




Fig. 3

4/15




Fig. 4




Fig. 5






Fig. 6

7/15

Increased uptake of folate-targeted imaging agent was seen in patient with inflamed joint

---




Fig. 7




Fig. 8 Folate-linked chelator EC20

9/15




Fig. 9



10/15




Fig. 10




Fig. 11

12/15




Fig. 12




Fig. 13






Fig. 14




Fig. 15

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 November 2002 (07.11.2002)

PCT

(10) International Publication Number  
**WO 02/087424 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 49/00**,  
39/395, 39/00, 39/385

(21) International Application Number: PCT/US02/13890

(22) International Filing Date: 2 May 2002 (02.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/288,208 2 May 2001 (02.05.2001) US

(71) Applicants and

(72) Inventors: **LOW, Philip, Stewart** [US/US]; 5850 Farm  
Ridge Road, West Lafayette, IN 47906 (US). **TURK,**  
**Mary, Jo** [US/US]; 1233 York Avenue, Apt. 13E, New  
York, NY 10026 (US).

(74) Agent: **LAMMERT, Steven, R.**; Barnes & Thornburg, 11  
South Meridian Street, Indianapolis, IN 46204 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
20 March 2003

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 02/087424 A3

(54) Title: TREATMENT AND DIAGNOSIS OF MACROPHAGE MEDIATED DISEASE

(57) Abstract: The invention relates to a method of treating or monitoring/diagnosing a disease state mediated by activated macrophages. The method comprises the step of administering to a patient suffering from a macrophage mediated disease state an effective amount of a composition comprising a conjugate or complex of the general formula: A<sub>b</sub>-X, where the group A<sub>b</sub> comprises a ligand capable of binding to activated macrophages, and when the conjugate is being used for treatment of the disease state, the group X comprises an immunogen, a cytotoxin, or a compound capable of altering macrophage function, and when the conjugate is being used for monitoring/diagnosing the disease state, X comprises an imaging agent. The method is useful for treating a patient suffering from a disease selected from the group consisting of rheumatoid arthritis, ulcerative colitis, Crohn's disease, inflammation, infections, osteomyelitis, atherosclerosis, organ transplant rejection, pulmonary fibrosis, sarcoidosis, and systemic sclerosis.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/13890

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 49/00,39/395, 39/00,39/385  
US CL : 424/9.1, 178.1, 184.1, 193.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 424/9.1, 178.1, 184.1, 193.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	WO 01/39806 A1 (ANTICANCER THERAPEUTIC INVENTIONS) 07 June 2001,(07.06.2001) see entire document	1-19
Y	WO 96/36367 A1 (PURDUE RESEARCH FOUNDATION) 21 November 1996 (21.11.96) see entire document.	1-19
Y	LU, J.Y. et al. Folate-Targeted Enzyme Prodrug Cancer Therapy Utilizing Penicillin-V Amidase and a Doxorubicin Prodrug. J. Drug Targeting. 1999, Vol. 7, No. 1, pages 43-53 (see entire document).	1-19
Y	LEAMON et al. Selective Targeting of Malignant Cells with Cytotoxin-Folate Conjugates. J. Drug Targeting. 1994, Vol. 2, No. 2, pages 101-112, (see entire document).	1-19
Y	SUDIMACK, J. et al. Targeted drug delivery via the folate receptor. Adv. Drug Deliv. Rev. March 2000, Vol. 41, No. 2 pages 147-162 (see entire document).	1-19
Y	REDDY et al. Folate-Mediated Targeting of Therapeutic and Imaging Agents to Cancers. Critical Reviews in Therapeutic Drug Carrier System. 1998, Vol. 15, pages 587-627 (see entire document).	1-19



Further documents are listed in the continuation of Box C.



See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;"

document member of the same patent family

Date of the actual completion of the international search

28 October 2002 (28.10.2002)

Date of mailing of the international search report

02 JAN 2003

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Michael A Belyavskyi

Telephone No. 703/308-0196

Form PCT/ISA/210 (second sheet) (July 1998)

# INTERNATIONAL SEARCH REPORT

PCT/US02/13890

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	REDDY, J. et al. Optimization of Folate-Conjugated Liposomal Vectors for Folate Receptor-Mediated Gene Therapy. J. Pharmaceutical Science. November 1999, Vol. 88, pages 1112-1118 (see entire document).	1-19

Form PCT/ISA/210 (second sheet) (July 1998)



# INTERNATIONAL SEARCH REPORT

PCT/US02/13890

## Continuation of B. FIELDS SEARCHED Item 3:

Biosis, Caplus, Medline, Embase, West, STN

Search terms: Low philip, folate receptor, binding ligand, cytotoxin, imaging agent, conjugates, liposones, antibody, macrophages.

Form PCT/ISA/210 (second sheet) (July 1998)